

# LA-UR-12-24196

Approved for public release; distribution is unlimited.

Title:	A discussion of numerical methods for capturing discontinuities in PDEs
Author(s):	Davis, Sean L.
Intended for:	Presentation



## Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



# Comparison of Methods for Capturing Discontinuities

With an Introduction to GPU Programming

Sean Davis and Will Matern



# Acknowledgements

- Mentor: Dr. Bob Robey
- Workshop Coordinator: Dr. Scott Runnels
- Collaborator: David Nicholaeff


Thank you!



# Outline

- Motivation
- Problems with capturing discontinuities
- Methods implemented and tested
  - First order Upwind
  - Lax-Wendroff
  - Limiters
  - MUSCL
  - SUPG
- Early MPDATA results
- GPU Programming
  - Introduction
  - Results solving the shallow water equations
- Conclusions

# Motivation



A simple model problem showing the advection of radioactive water dumped into the ocean and carried by a rotational current across the Pacific.

# 1D Advection


- We are using the 1D advection equation (in our 2D codes) to test various methods:

$$\frac{\partial \psi}{\partial t} + u \frac{\partial \psi}{\partial x} = 0$$

- We are looking at the advection of 3 different types of waves
  - Square Wave
  - Triangular Wave
  - Exponential Smooth Wave


# First Order Upwind Scheme

- Conservative 1<sup>st</sup> order scheme
- Overly dissipative



# Lax-Wendroff


- Conservative 2<sup>nd</sup> order scheme
- Approximate solution has oscillations around sharp gradients





# Limiters


- Godunov's Theorem – Any linear scheme above first order accurate (space) cannot be Total Variation Diminishing (TVD).



- Solution: Limiters (a nonlinear scheme)


# Lax-Wendroff w/ Superbee Limiter

- 1<sup>st</sup> order near discontinuities, 2<sup>nd</sup> order elsewhere
- For linear equations is mathematically guaranteed to be TVD



# Monotone Upstream Scheme for Conservation Laws (MUSCL)

- Used a modified Osher-Chakravarthy scheme
- Upwind biased
- Second order TVD approximation
- Numerical oscillations avoided using a minmod function



# Streamline Upwind Petrov-Galerkin (SUPG)


- Trial functions – a set of functions assumed to approximate the underlying solution. The trial functions have degrees of freedom that must be solved for.
- Test functions – a set of functions used to solve for the degrees of freedom of the trial functions. Uses the weak formulation of the governing equation.
- Petrov-Galerkin – Test functions  $\neq$  Trial functions
- SUPG – Choose test functions to be:

$$N + \frac{\alpha h}{2} \left( \frac{u}{\sqrt{u^2 + v^2}} \frac{\partial N}{\partial x} + \frac{v}{\sqrt{u^2 + v^2}} \frac{\partial N}{\partial y} \right)$$

where N is a trial function

# SUPG

- Inspired by finite difference methods (upwinding and artificial diffusion)
- Applies to linear advection/diffusion equations of any dimension
- Is neither TVD nor conservative




# Wave Advection Test Conclusions


- Lax-Wendroff with slope limiters
  - Effective for sharp discontinuities
  - Overly dissipative for thin waves (cuts off the top of the triangular/gaussian waves)
  - more dissipative than other methods
  - Superbee limiter artificially sharpens smooth waves
- MUSCL with ACM
  - most effective for problems with discontinuities
  - Unnaturally sharpens the solution
- SUPG
  - Balances well between diffusion and discontinuity sharpness
  - Doesn't introduce unnatural sharpening
  - Requires matrix operations
  - Non-conservative

# Results for 1D Advection in a 2D Code after 10 Periods


Lax-Wendroff (not TVD):




Lax-Wendroff (Superbee):



MUSCL:



SUPG:



# Multidimensional Positive Definite Advection Transport Algorithm (MPDATA)

- A donor-cell approximation is defined in terms of the local Courant number
- Adds a diffusive convective flux
- Subtract out added dissipation using an antidiffusive velocity
- As an example, the 1D advection equation,

$$\frac{\partial \Psi}{\partial t} = -\frac{\partial}{\partial x}(u\Psi)$$


- becomes,

$$\frac{\partial \Psi}{\partial t} = -\frac{\partial}{\partial x}(u\Psi) - \frac{\partial}{\partial x} \left( -K \frac{\partial \Psi}{\partial x} \right)$$



# MPDATA


- Antidiffusive correction to the donor cell method causes oscillations



PRELIMINARY

# MPDATA

- A Flux Corrected Transport (FCT) algorithm can correct the oscillations
- Performing additional local corrective iterations increases the order of accuracy of the solution




PRELIMINARY

# GPUs at a glance

- Much higher FLOPS (floating points operations per second) per dollar than CPUs.
- Different architecture from a CPU.
  - Many low throughput processing units
  - Cores share global memory
  - Warps/Wavefronts – groups of cores run same code.
- Must use a language compiled for GPU
  - OpenCL – Open programming language for multiple CPUs (on same node) and GPU.
  - CUDA – Nvidia only.
- This Project: OpenCL

# GPU architecture



<http://www.realworldtech.com/gt200/3/>

# Shallow Water Equations

- Simplification of the Euler Equations
  - Uniform Fluid Density
  - Wavelengths are much longer than fluid depth

$$\frac{\partial h}{\partial t} + \frac{\partial(uh)}{\partial x} + \frac{\partial(vh)}{\partial y} = 0$$


$$\frac{\partial(uh)}{\partial t} + \frac{\partial(u^2h + 1/2gh^2)}{\partial x} + \frac{\partial(uvh)}{\partial y} = 0$$

$$\frac{\partial(vh)}{\partial t} + \frac{\partial(uvh)}{\partial x} + \frac{\partial(v^2h + 1/2gh^2)}{\partial y} = 0$$


- Inert Tracer Equation

$$\frac{\partial \psi}{\partial t} + u \frac{\partial \psi}{\partial x} + v \frac{\partial \psi}{\partial y} = 0$$

# Shallow Water Simulations on the GPU



Fluid Height



Inert Tracer Concentration



# Future Directions

- FEM approach for full Euler equations
- Implement more of these schemes in the GPU code
- Develop full Euler Equation solvers with some of these schemes
- Alternative numerical methods (WENO, Discontinuous Galerkin, etc.)



# Question, Comments, Suggestions?